

11/09/00

JC965 U.S. PTO

11-14-80 A

Docket No. 62814-A/JPW/GJG

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Respectable Assistant Commissioner for Patents  
Washington, D.C. 20231

November 9, 2000

JC921 U.S. PTO  
09/711272  
11/09/00

RE:

Transmitted herewith for filing are the specification and claims of the patent application of: Timothy Norris, Jeffrey W. Raggon, Richard D. Connell, James D. Moyer, Michael J. Morin, Shana M. Kajiji, Barbara A. Foster, Karen J. Ferrante and

Inventor(s) Sandra L. Silberman for  
table Polymorph of N-(3-Ethynylphenylamino)-6, 7-Bis (2-Methoxyethoxy)-4-Quinazolinamine

Title of Invention Hydrochloride, Methods of Production, and Pharmaceutical Uses Thereof  
Also enclosed are:

☒ 4 sheet(s) of ☐ informal ☒ formal drawings.

☒ Oath or declaration of Applicant(s), unsigned

☒ A power of attorney, unsigned

☐ An assignment of the invention to \_\_\_\_\_

☒ A Preliminary Amendment

☐ A verified statement to establish small entity status under 37 C.F.R. §1.9 and §1.27.

The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

	NUMBER FILED		NUMBER EXTRA*		RATE		FEE	
					SMALL ENTITY	OTHER ENTITY	SMALL ENTITY	OTHER ENTITY
Total Claims	26 -20	=	6	X	\$ 9.00	\$ 18.00	= \$	\$ 108.00
Independent Claims	3 -3	=	0	X	\$ 40.00	\$ 80.00	= \$	\$ 0
Multiple Dependent Claims Presented: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No					\$ 135.00	\$ 270.00	= \$	\$
*If the different in Col. 1 is less than zero, enter "0" in Col. 2					BASIC FEE		\$ 355	\$ 710
					TOTAL FEE		\$	\$ 818.00

Applicants: Timothy Norris et al.  
Mail No.: Not Yet Known  
Filed: Herewith  
Number of Transmittals  
Page 2

- ☐ A check in the amount of \$ 818.00 to cover the filing fee.
- ☐ Please charge Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_.
- ☐ The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the following or credit any over-payment to Account No. 03-3125:
- ☒ Filing fees under 37 C.F.R. §1.16.
- ☒ Patent application processing fees under 37 C.F.R. §1.17.
- ☒ The issue fee set in 37 C.F.R. §1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. §1.311(b).
- ☐ Three copies of this sheet are enclosed.
- ☐ A certified copy of previously filed foreign application No. \_\_\_\_\_ filed in \_\_\_\_\_ on \_\_\_\_\_.
- Applicant(s) hereby claim priority based upon this aforementioned foreign application under 35 U.S.C. §119.
- ☒ Other (identify) Express Mail Certificate of Mailing dated  
November 9, 2000 and bearing Label No.  
EL 278 887 601 US.

Respectfully submitted,



John P. White  
Registration No. 28,678  
Attorney for Applicants  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10036  
(212) 278-0400

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

Applicants : Timothy Norris et al.  
Serial No. : Not Yet Known  
Filed : Herewith  
For : STABLE POLYMORPH OF N-(3-ETHYNYLPHENYL)-6,  
7-BIS (2-METHOXYETHOXY)-4-QUINAZOLINAMINE  
HYDROCHLORIDE, METHODS OF PRODUCTION, AND  
PHARMACEUTICAL USES THEREOF

1185 Avenue of the Americas  
New York, New York 10036  
November 9, 2000

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

**PRELIMINARY AMENDMENT**


Prior to examination, please make the following amendments to the  
above-identified application:

**In the Claims**

Please cancel claims 8-13, 33-49, and 51 without prejudice.

No fee, other than the enclosed application filing fee, is deemed  
necessary in connection with the filing of this Preliminary  
Amendment. However, if any fee is required, authorization is  
hereby given to charge the amount of any such fee to Deposit  
Account No. 03-3125.

Respectfully submitted,

  
John P. White  
Registration No. 28,678  
Gary J. Gershik  
Registration No. 39,992  
Attorneys for Applicants  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10036  
(212) 278-0400

*Application  
for  
United States Letters Patent*

To all whom it may concern:

Be it known that, we,  
Timothy Norris, Jeffrey W. Raggon, Richard D. Connell, James D. Moyer, Michael J. Morin, Shana M. Kajiji,  
Barbara A. Foster, Karen J. Ferrante and Sandra L. Silberman

have invented certain new and useful improvements in  
STABLE POLYMORPH OF N-(3-ETHYNYLPHENYLAMINO)-6, 7-BIS (2-METHOXYETHOXY)-4-  
QUINAZOLINAMINE HYDROCHLORIDE, METHODS OF PRODUCTION, AND PHARMACEUTICAL  
USES THEREOF

of which the following is a full, clear and exact description.

**STABLE POLYMORPH OF  
N-(3-ETHYNYLPHENYLAMINO)-6,7-BIS(2-  
5 METHOXYETHOXY)-4-QUINAZOLINAMINE HYDROCHLORIDE,  
METHODS OF PRODUCTION, AND PHARMACEUTICAL USES THEREOF**

This application claims the benefit of U.S. Provisional Application No. 60/206,420, filed May 23, 2000, U.S. Provisional  
10 Application No. 60/193,191, filed March 30, 2000, and U.S. Provisional Application No. 60/164,907, filed November 11, 1999, the contents of which are hereby incorporated by reference.

Throughout this application various publications are referenced.  
15 The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.


**20 Background of the Invention**

N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, in either its hydrochloride or mesylate forms, or in an anhydrous and hydrous form, is useful in the treatment of hyperproliferative disorders, such as cancers, in mammals.

25 U.S. Patent No. 5,747,498, issued May 5, 1998, which is incorporated herein by reference in its entirety, refers, in Example 20, to [6,7-bis(2-methoxyethoxy)-quinazolin-4-yl]-(3-ethynylphenyl)amine hydrochloride, which, the patent discloses,  
30 is an inhibitor of the erbB family of oncogenic and protooncogenic protein tyrosine kinases, such as epidermal growth factor receptor (EGFR), and is therefore useful for the treatment of proliferative disorders, such as cancers, in humans.

35 The mesylate form, described in PCT International Publication No. WO 99/55683 (PCT/IB99/00612, filed April 8, 1999), the entire disclosure of which is incorporated herein by reference,

and assigned to a common assignee, and shown in formula 1 below:



15 is useful for the treatment of proliferative disorders, and more preferred with parenteral methods of administration, as compared to the hydrochloride compound, i.e. with greater effectiveness in solution.

20 The mesylate compounds are more soluble in aqueous compositions than the hydrochloride compound, and thus the mesylate compounds are easily delivered according to parenteral methods of administration. The hydrochloride compound is however preferred with respect to solid administration such as with tablets and  
25 oral administration.

**Summary of the Invention**

The present invention relates to polymorphs, and methods for the selective production of polymorphs of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride, particularly in the stable polymorph form.

The present invention also relates to novel uses of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, in either its hydrochloride or mesylate forms, in an anhydrous or hydrous form, as well as in its various polymorph forms, in the treatment of hyperproliferative disorders, such as cancers, in mammals.

Description of the Figures

**Figure 1** The X-ray powder diffraction patterns for the hydrochloride polymorph A, the thermodynamically less stable form, over a larger range to show the first peaks.

**Figure 2** The X-ray powder diffraction patterns for the hydrochloride polymorph A, the thermodynamically less stable form, are over a shorter range to show more detail.

10

**Figure 3** The X-ray powder diffraction patterns for the hydrochloride polymorph B, the thermodynamically more stable form, over a larger range to show the first peaks.

15 **Figure 4** The X-ray powder diffraction patterns for the hydrochloride polymorph B, the thermodynamically more stable form, over a shorter range to show more detail.



**Detailed Description of the Invention**

Disclosed is a substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91. The polymorph is also characterized by the X-ray powder diffraction pattern shown in Figure 3.

10

Disclosed is a crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91, which is substantially free of the polymorph designated the A polymorph. The polymorph is also characterized by the X-ray powder diffraction pattern shown in Figure 3.

20

The polymorph designated the B polymorph may be in substantially pure form, relative to the A polymorph.

Also disclosed is a composition comprising a substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91. The hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine also exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately the values show in Table 3 or in Table 4 below.

And, the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the polymorph B form may be characterized by the X-ray powder diffraction pattern shown in Figure 3.

5

Also disclosed is a composition comprising a crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91 in a weight % of the B polymorph relative to the A polymorph which is at least 70%.. This composition may comprise at least 75% polymorph B, by weight; at least 80% polymorph B, by weight; at least 85% polymorph B, by weight; at least 90% polymorph B, by weight; at least 95% polymorph B, by weight; at least 97% polymorph B, by weight; at least 98% polymorph B, by weight; or at least 99% polymorph B, by weight relative to the A polymorph.

20

Further disclosed is a process for producing the polymorph B of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine by recrystallization of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in a solvent comprising alcohol and water.

25


In the process, the recrystallization may comprise the steps of:

- a) heating to reflux alcohol, water and the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine so as to form a solution;
- b) cooling the solution to between about 65 and 70 °C;
- c) clarifying the solution; and
- d) precipitating polymorph B by further cooling the clarified solution.


30

In the process, the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride is prepared by the steps of:


coupling a compound of formula 6



with a compound of formula 4




The compound of formula 6 is prepared by reacting a compound of formula 5



in a suspension of metal alkali and solvent and with heating.

The compound of formula 4 is prepared by chlorinating a compound of formula 3



Also disclosed is a pharmaceutical composition for the treatment of a hyperproliferative disorder in a mammal which substantially comprises a therapeutically effective amount of the polymorph B and a pharmaceutically acceptable carrier.

The pharmaceutical composition may be adapted for oral administration. It may be in the form of a tablet.

Also disclosed is a method of treating a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically effective amount of the polymorph B.

The method may be for the treatment of a cancer selected from brain, squamous cell, bladder, gastric, pancreatic, breast, head, neck, oesophageal, prostate, colorectal, lung, renal, kidney, ovarian, gynecological and thyroid cancer.

The method may also be for the treatment of a cancer selected from non-small cell lung cancer (NSCLC), refractory ovarian cancer, head and neck cancer, colorectal cancer and renal cancer.

In the method, the therapeutically effective amount may be from about 0.001 to about 100 mg/kg/day, or from about 1 to about 35 mg/kg/day.

In the method, the therapeutically effective amount may also be


from about 1 to about 7000 mg/day; from about 5 to about 2500 mg/day; from about 5 to about 200 mg/day; or from about 25 to about 200 mg/day.

- 5 Further disclosed is a method for the treatment of a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically effective amount of the polymorph B in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, 10 alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, and anti-androgens.
- 15 Yet further disclosed is a method of making a composition which composition comprises substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having 20 characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91, comprising admixing the crystalline polymorph designated the B polymorph with a carrier.
- 25 The carrier may be a pharmaceutically acceptable carrier.

Also disclosed is a method of preparing polymorph B of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride which comprises the step of recrystallizing N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in a solvent comprising alcohol.


In the method the solvent may further comprises water.

In the method, the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride is prepared by coupling a compound of formula 6




6

with a compound of formula 4



4.


In the method, the compound of formula 6 is prepared by reacting a compound of formula 5



5

in a suspension of metal alkali and solvent and with heating.


In the method, the compound of formula 4 is prepared by chlorinating a compound of formula 3




3.

Further disclosed is a method for the production of the polymorph B of claim 1 comprising the steps of:

- a) substitution chlorination of starting quinazolinamine  
15 compound of formula 3



having an hydroxyl group, to provide a compound of formula 4




4

by reaction thereof in a solvent mixture of thionyl chloride,  
35 methylene chloride and dimethylformamide,

b) preparation of a compound of formula 6


5



6

10 in situ from starting material of compound of formula 5

15



5

20

by reaction of the latter in a suspension of metal alkali and solvent and with heating;

c) reaction of the compound of formula 6 in situ with the  
25 compound of formula 4 wherein the compound of formula 6 replaces the chlorine in the compound of formula 4 to give the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride;

d) recrystallizing the N-(3-ethynylphenyl)-6,7-bis(2-  
30 methoxyethoxy)-4-quinazolinamine hydrochloride, in alcohol, into the polymorph B form.

In this method, the substitution chlorination may be quenched in the presence of aqueous sodium hydroxide; aqueous sodium  
35 bicarbonate; aqueous potassium hydroxide; aqueous potassium bicarbonate; aqueous potassium carbonate; aqueous sodium



carbonate, or a mixture thereof.

Yet further disclosed is a method for the production of polymorph B of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine by recrystallization comprising the steps of:

- a) heating to reflux alcohol, water and the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine so as to form a solution;
- b) cooling the solution to between about 65 and 70 °C;
- c) clarifying the solution; and
- d) precipitating polymorph B by further cooling the clarified solution.

Further disclosed is a composition consisting essentially of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the form of polymorph A, which is characterized by the following peaks in its X-ray powder diffraction pattern shown in Figure 1.

Also disclosed is a composition consisting essentially of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the form of polymorph A, which is characterized by the peaks shown in Table 1 or Table 2 below.

A prodrug of any of the compound herein is also disclosed.

Further disclosed is a method of inducing differentiation of tumor cells in a tumor comprising contacting the cells with an effective amount of any of the compounds or compositions disclosed herein.

Also disclosed is a method for the treatment of NSCLC (non small cell lung cancer), pediatric malignancies, cervical and other tumors caused or promoted by human papilloma virus (HPV),

melanoma, Barrett's esophagus (pre-malignant syndrome), adrenal and skin cancers and auto immune, neoplastic cutaneous diseases and atherosclerosis in a mammal comprising administering to said mammal a therapeutically effective amount of a pharmaceutical composition comprised of at least one of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and pharmaceutically acceptable salts thereof in anhydrous and hydrate forms.

The treatment may further comprise a palliative or neo-  
10 adjuvant/adjuvant monotherapy; or comprises blocking epidermal growth factor receptors (EGFR).

The method of may also be used in the treatment of tumors that express EGFRVIII.

15 The method may further comprise a combination with any of chemotherapy and immunotherapy; or treatment with either or both anti-EGFR and anti-EGF antibodies; or administration to said mammal of a member of the group consisting of inhibitors of MMP  
20 (matrix-metallo-proteinase), VEGFR (vascular endothelial growth factor receptor), farnesyl transferase, CTLA<sub>4</sub> (cytotoxic T-lymphocyte antigen 4) and erbB2, MAb to VEGFr, rhuMAb-VEGF, erbB2 MAb and avb3 Mab.


25 The pharmaceutical compounds used may be radiation sensitizers for cancer treatment or in combination with anti-hormonal therapies, or for the inhibition of tumor growth in humans in a regimen with radiation treatment.

30 Further disclosed is a method for the chemoprevention of basal or squamous cell carcinoma of the skin in areas exposed to the sun or in persons of high risk to said carcinoma, said method comprising administering to said persons a therapeutically effective amount of a pharmaceutical composition comprised of at least one of N-(3-  
35 ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and

pharmaceutically acceptable salts thereof in anhydrous and hydrate forms.

Also is a method of inducing differentiation of tumor cells in a tumor comprising contacting the cells with an effective amount of the compound of at least one of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and pharmaceutically acceptable salts thereof in anhydrous and hydrate forms.

It is accordingly an object of the present invention to provide a method for the production of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine in HCl form (Formula 2):



## 2 (Polymorph form A and B)

making it more suitable for tablet and oral administration and consisting essentially of the stable polymorphic form (polymorph form B) as well as the compound in such polymorph B form and the intermediate polymorph A in essentially pure form.

It is a further object of the present invention to provide such stable polymorph form B in a pharmaceutical orally administered composition.

Stability of the hydrochloride compound is of concern for its use in the treatment of patients since variations will affect effective dosage level and administration. It has been discovered that the hydrochloride of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine exists in two polymorph states, polymorph A and B. This contrasts with the mesylate compounds which exist in three polymorph states (mesylate polymorphs A, B and C). Polymorph B of the hydrochloride was found to be the thermodynamically most stable and desirable form and the present invention comprises the polymorph B compound in the substantially pure polymorphic B form and pharmaceutical compositions of the substantially pure form of polymorph B, particularly in tablet form and a method of the selective production of the compound.

15 The hydrochloride compound disclosed in the U.S. Patent No. 5,747,498 actually comprised a mixture of the polymorphs A and B, which, because of its partially reduced stability (i.e., from the polymorph A component) was not more preferred for tablet form than the mesylate salt forms.

20 Specifically, the present invention relates to methods of producing the hydrochloride compound forms of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine and for producing the stable form B in high yield. The mesylate salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine has been discovered to exist in at least three polymorphic forms which have been designated A, B, and C, of increasing stability with different X-ray powder diffraction patterns. The X-ray powder diffraction patterns for the hydrochloride polymorph A ( $A_1$  and  $A_2$ ) and B ( $B_1$  and  $B_2$ ) forms are shown in Figures 1-4 as follows: graphs of Figures 1 and 3 are over a larger range to fully show the first peaks for A and B, respectively, and graphs of Figures 2 and 4 are over a shorter range to show more overall detail for A and B, respectively.

The data contained in the above X-ray diffraction patterns of Figure 1-4 are tabulated in the following Tables 1-4:

Table:1 Polymorph A

Anode: Cu- Wavelength 1: 1.54056 Wavelength 2: 1.54439 (Rel Intensity: 0.500)

5 Range# 1 -Coupled: 3.000 to 40.000 StepSize: 0.040 StepTime: 1.00

Smoothing Width: 0.300 Threshold: 1.0

d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)
15.82794	100.0	6.63179	1.7	4.54453	4.8	3.61674	8.2	2.91238	3.5
14.32371	3.9	5.84901	2.1	4.19685	4.7	3.50393	9.3	2.73148	3.7
11.74376	1.5	5.69971	2.3	4.16411	4.4	3.40200	6.0	2.60193	1.8
11.03408	1.2	5.46922	2.4	3.97273	4.7	3.35174	5.3	2.48243	1.3
10.16026	1.4	5.21396	3.6	3.91344	12.4	3.29005	4.2	2.40227	2.2
8.98039	13.1	4.80569	3.5	3.78223	24.2	3.05178	7.1	2.31297	1.7
7.85825	7.8	4.70077	12.2	3.67845	8.8	2.97750	3.0		

Table:2 Polymorph A

Anode: Cu - Wavelength 1: 1.54056 Wavelength 2: 1.54439 (Rel Intensity: 0.500)

Range#1 - Coupled: 3.000 to 40.000 StepSize: 0.040 StepTime: 1.00

20 Smoothing Width: 0.300 Threshold: 1.0

2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)
5.579	100.0	13.340	1.7	19.517	4.8	24.594	8.2	30.673	3.5
6.165	3.9	15.135	2.1	21.152	4.7	25.398	9.3	32.759	3.7
7.522	1.5	15.534	2.3	21.320	4.4	26.173	6.0	34.440	1.8
8.006	1.2	16.193	2.4	22.360	4.7	26.572	5.3	36.154	1.3
8.696	1.4	16.991	3.6	22.703	12.4	27.080	4.2	37.404	2.2
9.841	13.1	18.447	3.5	23.502	24.2	29.240	7.1	38.905	1.7
11.251	7.8	18.862	12.2	24.175	8.8	30.007	3.0		

Table:3 Polymorph B

Anode: Cu - Wavelength 1 1.54056 Wavelength 2: 1.54439 (Rel Intensity:0.500)

Range # 1 - Coupled 3.000 to. 40.040 StepSize: 0.040 StepTime 1.00

Smoothing Width: 0.300 Threshold: 1.0

d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)
14.11826	100.0	5.01567	2.5	3.86656	4.8	3.23688	0.9	2.74020	1.7
11.23947	3.2	4.87215	0.7	3.76849	2.3	3.16755	1.5	2.69265	1.7
9.25019	3.9	4.72882	1.5	3.71927	3.0	3.11673	4.3	2.58169	1.5
7.74623	1.5	4.57666	1.0	3.63632	6.8	3.07644	1.4	2.51043	0.8
7.08519	6.4	4.39330	14.4	3.53967	10.0	2.99596	2.1	2.47356	1.0
6.60941	9.6	4.28038	4.2	3.47448	3.7	2.95049	0.9	2.43974	0.6
5.98828	2.1	4.20645	14.4	3.43610	3.9	2.89151	1.6	2.41068	1.1
5.63253	2.9	4.06007	4.7	3.35732	2.8	2.83992	2.2	2.38755	1.4
5.22369	5.5	3.95667	4.5	3.31029	5.6	2.81037	2.4	2.35914	1.7

Table:4 Polymorph B

Anode: Cu - Wavelength 1 1.54056 Wavelength 2: 1.54439 (Rel Intensity:0.500)

Range# 1 - Coupled: 3.000 to 40.040 StepSize 0.040 StepTime: 1.00

Smoothing Width:0.300 Threshold: 1.0

5	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)
	6.255	100.0	17.668	2.5	22.982	4.8	27.534	0.9	32.652	1.7
	7.860	3.2	18.193	0.7	23.589	2.3	28.148	1.5	33.245	1.7
	9.553	3.9	18.749	1.5	23.906	3.0	28.617	4.3	34.719	1.5
	11.414	1.5	19.379	1.0	24.459	6.8	29.000	1.4	35.737	0.8
10	12.483	6.4	20.196	14.4	25.138	10.0	29.797	2.1	36.288	1.0
	13.385	9.6	20.734	4.2	25.617	3.7	30.267	0.9	36.809	0.6
	14.781	2.1	21.103	14.4	25.908	3.9	30.900	1.6	37.269	1.1
	15.720	2.9	21.873	4.7	26.527	2.8	31.475	2.2	37.643	1.4
	16.959	5.5	22.452	4.5	26.911	5.6	31.815	2.4	38.114	1.7

15

It is to be understood that the X-ray powder diffraction pattern is only one of many ways to characterize the arrangement of atoms comprising the compound N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride, and that other methods well known in the art, e.g. single crystal X-ray diffraction, may be used to identify in a sample, composition or other preparation the presence of polymorph B of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine.

25

The present invention relates to a compound which is polymorph B of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91. This invention also relates to a polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately the values shown in Table 4 above.

35

This invention also relates to a compound which is polymorph A of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-

methoxyethoxy)-4-quinazolinamine that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 5.58, 9.84, 11.25, 18.86, 22.70, 23.50, 24.18, 24.59, 25.40 and 29.24. This invention also relates to a polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately the values shown in Table 2 above.

10


#### Method of Production

The polymorph B in substantially pure form of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride (compound of formula 1) is prepared, in accordance with the method of the present invention, by the steps of;

15

1) substitution chlorination of starting quinazolinamine compound (formula 3):

20




3

25


having an hydroxyl group, such as by reaction thereof in a solvent mixture of thionyl chloride, methylene chloride, and dimethylformamide, and finally quenching the reaction with an aqueous solution of sodium hydroxide or sodium bicarbonate. The compound of formula 4:

30




is produced in high yield with replacement of the hydroxyl group with chlorine;

2) preparation of compound of formula 6:



from starting material of formula 5:



by reaction of the latter in a suspension of NaOH (or KOH, or a combination) in toluene with heating;



3) reaction of the compound of formula 6 with the compound of formula 4 of step 1 wherein the compound of formula 6 replaces the chlorine to give the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride (compound of formula 2) with a 97%  
5 yield.

4) recrystallization of the compound of formula 2 (comprising both polymorph A and polymorph B) into the more stable polymorph B in a solvent comprising alcohol (e.g. 2B-ethanol) and water,  
10 generally in high yield, e.g., about 85%.


Accordingly, the present invention relates to a method of preparing polymorph B of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride which comprises  
15 recrystallization of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in a solvent comprising alcohol and water. In one embodiment, the method comprises the steps of heating to reflux alcohol, water and the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine so as  
20 to form a solution; cooling the solution to between about 65 and 70 °C; clarifying the solution; and precipitating polymorph B by further cooling the clarified solution. In an embodiment, the alcohol is ethanol. In a preferred embodiment, the ratio of ethanol to water is about 4:1. It is to be expected that other  
25 lower alcohols, e.g., C<sub>1</sub>-C<sub>4</sub> alcohols, are also suitable for recrystallization of polymorph B with adjustment of the alcohol to water ratio as needed. In another preferred embodiment, the compound to be recrystallized is present in an amount relative to the total volume of solvent at a weight to volume ratio of about  
30 0.05. In an embodiment, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride is prepared by coupling a compound of formula 6 with a compound of formula 4. In another embodiment, the compound of formula 6 is prepared by reacting a compound of formula 5 in a suspension of metal alkali  
35 and solvent, with heating.

In an embodiment, the compound of formula 4 is prepared by chlorinating a compound of formula 3 by reaction of the latter in a solvent mixture of thionyl chloride, methylene chloride and dimethylformide, and subsequently quenching the reaction with an aqueous solution of sodium hydroxide. Alternatively, an aqueous solution of sodium bicarbonate can be substituted for the sodium hydroxide solution.

This invention relates to polymorph B of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine prepared by the above methods. In an embodiment, the polymorph B is prepared by using the starting materials described herein. In a preferred embodiment, polymorph B is prepared by reaction of the starting materials described herein with the reagents and conditions according to the methods described herein and in the Examples which follow.

### General synthesis

N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride has been found to exist in two distinct anhydrous polymorphic forms A and B. The production method for the various polymorphs is with components separately reacted in accordance with the following scheme:



### Uses

As described in the aforementioned U.S. Patent No. 5,747,498 and PCT International Publication No. WO 99/55683, the compounds made in accordance with the present invention are useful for the  
5 treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of the hydrochloride or mesylate form of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and a pharmaceutically acceptable carrier.

- 10 The term "compound(s) of the invention" referred to herein is preferably the polymorph B form of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride, but is not meant to exclude the mesylate form and its three polymorphs, or polymorph A of the hydrochloride form, or  
15 a mixture of polymorphs B and A of the hydrochloride form or other non-crystalline forms of the compound.

- The term "treating" as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or  
20 preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, refers to the act of treating, as "treating" is defined immediately above.

- 25 "Abnormal cell growth", as used herein, refers to cell growth that is independent of normal regulatory mechanisms (e.g., loss of contact inhibition), including the abnormal growth of normal cells and the growth of abnormal cells. This includes, but is not limited to, the abnormal growth of: (1) tumor cells (tumors), both  
30 benign and malignant, expressing an activated Ras oncogene; (2) tumor cells, both benign and malignant, in which the Ras protein is activated as a result of oncogenic mutation in another gene; (3) benign and malignant cells of other proliferative diseases in which aberrant Ras activation occurs. Examples of such benign  
35 proliferative diseases are psoriasis, benign prostatic

hypertrophy, human papilloma virus (HPV), and restenosis. "Abnormal cell growth" as used herein also refers to and includes the abnormal growth of cells, both benign and malignant, resulting from activity of the enzymes farnesyl protein transferase, protein  
5 kinases, protein phosphatases, lipid kinases, lipid phosphatases, or activity of transcription factors, or intracellular or cell surface receptor proteins.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine  
10 hydrochloride, preferably the stable polymorph B form, is additionally used for the treatment of a variety of additional human tumors containing hyperproliferating cells that are activated by the signal transduction pathways stimulated by EGFR, whether by overexpression (e.g. due to one or more of - altered  
15 transcription, altered mRNA degradation or gene amplification) of the EGFR protein itself, another receptor protein with which EGFR can form active heterodimers, or one of the ligands that activate EGFR (e.g. EGF, TGF $\alpha$ , amphiregulin,  $\beta$ -cellulin, heparin-binding EGF, or epiregulin) or a heterodimerizing receptor, or due to a  
20 dependence or partial dependence on the activity of a "normal" level of EGFR protein, whether activated by extracellular ligand, intracellular signal transduction pathways and/or genetic alterations or polymorphisms that result in amino acid substitutions that produce increased or ligand-independent  
25 activity (e.g. EGFRvIII, Archer G.E. et. al. (1999) Clinical Cancer Research 5:2646-2652). Such tumors, including both benign and malignant, include renal (such as kidney, renal cell carcinoma, or carcinoma of the renal pelvis), liver, kidney, bladder (particularly invasive tumors), breast (including estrogen  
30 receptor negative and positive tumors, and progesterone receptor negative and positive tumors), gastric, esophageal (including Barrett's mucosa, squamous cell carcinomas and adenocarcinomas), larynx, ovarian, colorectal (particularly deeply invasive tumors), including anal, prostate, pancreatic, lung (particularly non-small  
35 cell lung cancer (NSCLC) adenocarcinomas, large cell tumors and

squamous cell carcinomas, but also reactive (squamous metaplasia and inflammatory atypia) as well as precancerous (dysplasia and carcinoma in situ) bronchial lesions associated with both NSCLC adenocarcinomas and squamous cell carcinomas), gynecological, including vulval, endometrial, uterine (e.g, sarcomas), cervical, vaginal, vulval, and fallopian tube cancers, thyroid, hepatic carcinomas, skin cancers, sarcomas, brain tumors, including glioblastomas (including glioblastoma multiforme), astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas and pituitary adenomas, and various other head and neck tumors (particularly squamous cell carcinomas), and metastases of all of the above.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride, preferably the stable polymorph B form, is also used for the treatment of a variety of additional human hyperplastic conditions containing hyperproliferating cells that are activated by the signal transduction pathways capable of stimulation by EGFR, such as benign hyperplasia of the skin (e.g. psoriasis) or prostate (e.g. BPH), chronic pancreatitis, or reactive hyperplasia of pancreatic ductal epithelium, or kidney disease (including proliferative glomerulonephritis and diabetes-induced renal disease) in a mammal which composition comprises a therapeutically effective amount of the hydrochloride of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, preferably the polymorph B form, and a pharmaceutically acceptable carrier.

In addition, pharmaceutical compositions including the compounds made in accordance with the present invention provide for the prevention of blastocyte implantation in a mammal, which composition comprises a therapeutically effective amount of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride, preferably the polymorph B form, and a pharmaceutically acceptable carrier.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride, preferably the stable polymorph B form, is also used for the treatment of additional disorders in which cells are activated by the signal transduction pathways stimulated by EGFR, whether by overexpression (due to one or more of - altered transcription, altered mRNA degradation or gene amplification) of the EGFR protein itself, another receptor protein with which it can form active heterodimers, or one of the ligands that activate EGFR (e.g. EGF, TGF $\alpha$ , amphiregulin,  $\beta$ -cellulin, heparin-binding EGF, or epiregulin) or a heterodimerizing receptor, or due to a dependence or partial dependence on the activity of a "normal" level of EGFR protein, whether activated by extracellular ligand, intracellular signal transduction pathways and/or genetic alterations or polymorphisms that result in amino acid substitutions that produce increased or ligand-independent activity (e.g. EGFRvIII, Archer G.E. et. al. (1999) Clinical Cancer Research 5:2646-2652). Such disorders may include those of a neuronal, glial, astrocytal, hypothalamic, and other glandular, macrophagal, epithelial, stromal, or blastocoelic nature in which aberrant or 'normal' function, expression, activation or signalling via EGFR may be involved. Such disorders may furthermore involve the modulation by EGF (or other ligands that activate EGFR or heterodimerizing receptors) of adipocyte lipogenesis, bone resorption, hypothalamic CRH release, hepatic fat accumulation, T-cell proliferation, skin tissue proliferation or differentiation, corneal epithelial tissue proliferation or differentiation, macrophage chemotaxis or phagocytosis, astroglial proliferation, wound healing, polycystic kidney disease, lung epithelial proliferation or differentiation (e.g. associated with asthmatic airway remodeling or tissue repair), inflammatory arthritis (e.g. rheumatoid arthritis, systemic lupus erythematosus-associated arthritis, psoriatis arthritis) testicular androgen production, thymic epithelial cell proliferation, uterine epithelial cell proliferation, angiogenesis, cell survival, apoptosis, NF $\kappa$ B activation, vascular

smooth muscle cell proliferation, restenosis or lung liquid secretion.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine  
5 hydrochloride, preferably the stable polymorph B form, is also  
used for the treatment of a range of leukemias (chronic and acute)  
and lymphoid malignancies (e.g. lymphocytic lymphomas), diabetes,  
diabetic and other retinopathies, such as retinopathy or  
prematurity, age-related macular degeneration, solid tumors of  
10 childhood, glioma, hemangiomas, melanomas, including intraocular  
or uveal melanomas, Kaposi's sarcoma, Hodgkin's disease,  
epidermoid cancers, cancers of the endocrine system (e.g.  
parathyroid, adrenal glands), bone small intestine, urethra, penis  
and ureter, atherosclerosis, skin diseases such as eczema and  
15 scleroderma, mycoses fungoides, sarcomas of the soft tissues and  
neoplasm of the central nervous system (e.g. primary CNS lymphoma,  
spinal axis tumors, brain stem gliomas, or pituitary adenomas).

The treatment of any of the hyperproliferative or additional  
20 disorders described above may be applied as a monotherapy, or may  
involve in addition to [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-  
(3-ethynylphenyl)-amine hydrochloride, preferably the stable  
polymorph B form, application with one or more additional drugs or  
treatments (e.g. radiotherapy, chemoradiotherapy) that are anti-  
25 hyperproliferative, anti-tumor or antihyperplastic in nature.  
Such conjoint treatment may be achieved by way of simultaneous,  
sequential, cyclic or separate dosing of the individual components  
of the treatment. [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-  
ethynylphenyl)-amine hydrochloride, preferably the stable  
30 polymorph B form, is typically used at doses of 1-7000 mg/day,  
preferably 5-2500 mg/day, most preferably 5-200 mg/day, for any of  
the above treatments.

Furthermore, the various forms of N-(3-ethynylphenyl)-6,7-bis(2-  
35 methoxyethoxy)-4-quinazolinamine including the mesylate and

hydrochloride forms (all polymorph forms) as well as other pharmaceutically acceptable salt forms, and anhydrous and hydrate forms, can be used for treatment, with a therapeutically-effective amount of the aforementioned compounds and a pharmaceutically acceptable carrier, of the specific conditions of NSCLC (non small cell lung cancer), pediatric malignancies, cervical and other tumors caused or promoted by human papilloma virus (HPV), melanoma, Barrett's esophagus (pre-malignant syndrome) and adrenal and skin cancers as well as auto immune and neoplastic cutaneous diseases such as mycoses fungoides, in a mammal, as well as for the chemoprevention of basal or squamous cell carcinomas of the skin, especially in areas exposed to the sun or in persons known to be at high risk for such cancers. In addition, the aforementioned compounds are useful in treatment of atherosclerosis, with epidermal growth factor having been implicated in the hyperproliferation of vascular smooth muscle cells responsible for atherosclerotic plaques (G.E. Peoples et al., Proc. Nat. Acad. Sci. USA 92:6547-6551, 1995).

The compounds of the present invention are potent inhibitors of the erbB family of oncogenic and protooncogenic protein tyrosine kinases such as epidermal growth factor receptor (EGFR), *erbB2*, *HER3*, or *HER4* and thus are all adapted to therapeutic use as antiproliferative agents (e.g., anticancer) in mammals, particularly in humans. The compounds of the present invention are also inhibitors of angiogenesis and/or vasculogenesis.

The compounds of the present invention may also be useful in the treatment of additional disorders in which aberrant expression of ligand/receptor interactions or activation or signalling events related to various protein tyrosine kinases are involved. Such disorders may include those of neuronal, glial, astrocytal, hypothalamic, glandular, macrophagal, epithelial, stromal, or blastocoelic nature in which aberrant function, expression, activation or signalling of the erbB tyrosine kinases are



involved. In addition, the compounds of the present invention may have therapeutic utility in inflammatory, angiogenic and immunologic disorders involving both identified and as yet unidentified tyrosine kinases that are inhibited by the compounds  
5 of the present invention.

In addition to direct treatment of the above ailments with the compounds, the utilization and treatment in these and general applications may be as palliative or neo-adjuvant/adjuvant  
10 monotherapy, in blocking epidermal growth factor receptors (EGFR) and for use in treatment of tumors that express a variant form of EGFR known as EGFRvIII as described in the scientific literature (e.g., DK Moscatello et al. Cancer Res. 55:5536-5539, 1995), as well as in a combination with chemotherapy and immunotherapy. As  
15 described in more detail below, treatment is also possible with both anti-EGFR and anti-EGF antibody combinations or with combination of inhibitors of MMP (matrix-metallo-proteinase), other tyrosine kinases including VEGFR (vascular endothelial growth factor receptor), farnesyl transferase, CTLA<sub>4</sub> (cytotoxic  
20 T-lymphocyte antigen 4) and erbB2. Further treatments include MAb to VEGFR, and other cancer-related antibodies including rhuMAb-VEGF (Genentech, Phase III), the erbB2 MAb available as Herceptin (Genentech, Phase III), or the avb3 MAb available as Vitaxin (Applied Molecular Evolution/MedImmune, Phase II).

25

The invention also relates to a pharmaceutical composition and a method of treating any of the mentioned disorders in a mammal which comprises administering to said mammal a therapeutically effective amount of N-(3-ethynylphenyl)-6,7-bis(2-  
30 methoxyethoxy)-4-quinazolinamine, preferably in hydrochloride polymorph B form, and a pharmaceutically acceptable carrier.

### Combination Therapy

The active compound may be applied as a sole therapy or may involve one or more other materials and treatment agents such as both anti-EGFR and anti-EGF antibody combinations or with  
5 combination of inhibitors of MMP (matrix-metallo-proteinase), other tyrosine kinases including VEGFR (vascular endothelial growth factor receptor), farnesyl transferase, CTLA<sub>4</sub> (cytotoxic T-lymphocyte antigen 4) and erbB2, as well as MAb to VEGFr, and other cancer-related antibodies including rhuMAb-VEGF, the erbB2  
10 MAb, or avb3.

Thus, the active compound may be applied with one or more other anti-tumor substances, for example those selected from, for example, mitotic inhibitors, for example vinblastine; alkylating  
15 agents, for example cis-platin, carboplatin and cyclophosphamide; anti-metabolites, for example 5-fluorouracil, cytosine arabinoside and hydroxyurea, or, for example, one of the preferred anti-metabolites disclosed in European Patent Application No. 239362 such as N-(5-[N-(3,4-dihydro-2-methyl-4-  
20 oxoquinazolin-6-ylmethyl)-N-methylamino]-2-thenoyl)-L-glutamic acid; growth factor inhibitors; cell cycle inhibitors; intercalating antibiotics, for example adriamycin and bleomycin; enzymes, for example interferon; and anti-hormones, for example anti-estrogens such as Nolvadex® (tamoxifen) or, for example  
25 anti-androgens such as Casodex® (4'-cyano-3-(4-fluorophenylsulphonyl)-2-hydroxy-2-methyl-3'-(trifluoromethyl)propionanilide).

In a further embodiment, the compounds of the invention may be  
30 administered in conjunction with an anti-angiogenesis agent(s) such as a MMP-2 (matrix-metalloproteinase-2) inhibitor(s), a MMP-9 (matrix-metalloproteinase-9) inhibitor(s), and/or COX-II (cyclooxygenase II) inhibitor(s) in the methods of treatment and compositions described herein. For the combination therapies  
35 and pharmaceutical compositions described herein, the effective

amounts of the compound of the invention and of the chemotherapeutic or other agent useful for inhibiting abnormal cell growth (e.g., other antiproliferative agent, anti-angiogenic, signal transduction inhibitor or immune-system enhancer) can be determined by those of ordinary skill in the art, based on the effective amounts for the compound described herein and those known or described for the chemotherapeutic or other agent. The formulations and routes of administration for such therapies and compositions can be based on the information described herein for compositions and therapies comprising the compound of the invention as the sole active agent and on information provided for the chemotherapeutic or other agent in combination therewith.

15 The invention also relates to production of compounds used in a method for the treatment of a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically effective amount of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, and anti-androgens.

25

The compounds are also useful as radiation sensitizers for cancer treatment and may be combined with anti-hormonal therapies. Parameters of adjuvant radiation therapies are for example contained in PCT/US99/10741, as published on 25 November 1999, in International Publication No. WO 99/60023, the disclosure of which is included herein by reference thereto. With such mode of treatment for example, for inhibiting tumor growth, a radiation dosage of 1-100 Gy is utilized preferably in conjunction with at least 50 mg of the pharmaceutical compound, in a preferred dosage regimen of at least five days a week for about two to ten weeks.

Thus, this invention further relates to a method for inhibiting abnormal cell growth in a mammal which method comprises administering to the mammal an amount of the compound of the invention, or a pharmaceutically acceptable salt or solvate or  
5 prodrug thereof, in combination with radiation therapy, wherein the amount of the compound, salt, solvate or prodrug is in combination with the radiation therapy effective in inhibiting abnormal cell growth in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can  
10 be used in the combination therapy described herein.

Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-II (cyclooxygenase II) inhibitors, can be used in conjunction  
15 with the compound of the invention in the methods and pharmaceutical compositions described herein. Examples of useful COX-II inhibitors include CELEBREX™ (alecoxib), valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published October 24, 1996), WO  
20 96/27583 (published March 7, 1996), European Patent Application No. 97304971.1 (filed July 8, 1997), European Patent Application No. 99308617.2 (filed October 29, 1999), WO 98/07697 (published February 26, 1998), WO 98/03516 (published January 29, 1998), WO 98/34918 (published August 13, 1998), WO 98/34915 (published  
25 August 13, 1998), WO 98/33768 (published August 6, 1998), WO 98/30566 (published July 16, 1998), European Patent Publication 606,046 (published July 13, 1994), European Patent Publication 931,788 (published July 28, 1999), WO 90/05719 (published May 31, 1990), WO 99/52910 (published October 21, 1999), WO 99/52889  
30 (published October 21, 1999), WO 99/29667 (published June 17, 1999), PCT International Application No. PCT/IB98/01113 (filed July 21, 1998), European Patent Application No. 99302232.1 (filed March 25, 1999), Great Britain patent application number 9912961.1 (filed June 3, 1999), United States Provisional Application No.  
35 60/148,464 (filed August 12, 1999), United States Patent 5,863,949

(issued January 26, 1999), United States Patent 5,861,510 (issued January 19, 1999), and European Patent Publication 780,386 (published June 25, 1997), all of which are incorporated herein in their entireties by reference. Preferred MMP-2 and MMP-9 inhibitors are those that have little or no activity inhibiting MMP-1. More preferred, are those that selectively inhibit MMP-2 and/or MMP-9 relative to the other matrix-metalloproteinases (*i.e.* MMP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13).

10

Some specific examples of MMP inhibitors useful in the present invention are AG-3340, RO 32-3555, RS 13-0830, and the compounds recited in the following list:

15 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclopentyl)-amino]-propionic acid;

3-exo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide;

20 (2R, 3R) 1-[4-(2-chloro-4-fluoro-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide;

4-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide;

25

3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclobutyl)-amino]-propionic acid;

30 4-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide;

(R) 3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-3-carboxylic acid hydroxyamide;

35 (2R, 3R) 1-[4-(4-fluoro-2-methyl-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide;

3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-1-methyl-ethyl)-amino]-propionic acid;

40

3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(4-hydroxycarbamoyl-tetrahydro-pyran-4-yl)-amino]-propionic acid;

45 3-exo-3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide;

3-endo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; and

(R) 3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-  
5 furan-3-carboxylic acid hydroxyamide;

and pharmaceutically acceptable salts and solvates of said compounds.

10 Other anti-angiogenesis agents, including other COX-II inhibitors and other MMP inhibitors, can also be used in the present invention.

The compound of the present invention can also be used with signal  
15 transduction inhibitors, such as other agents that can inhibit EGFR (epidermal growth factor receptor) responses, such as EGFR antibodies, EGF antibodies, and other molecules that are EGFR inhibitors; VEGF (vascular endothelial growth factor) inhibitors, such as VEGF receptors and molecules that can inhibit VEGF; and  
20 erbB2 receptor inhibitors, such as other organic molecules or antibodies that bind to the erbB2 receptor, for example, HERCEPTIN™ (Genentech, Inc. of South San Francisco, California, USA).

25 EGFR inhibitors are described in, for example in WO 95/19970 (published July 27, 1995), WO 98/14451 (published April 9, 1998), WO 98/02434 (published January 22, 1998), and other compounds described in United States Patent 5,747,498 (issued May 5, 1998), and such substances can be used in the present invention as  
30 described herein. EGFR-inhibiting agents include, but are not limited to, the monoclonal antibodies C225 and anti-EGFR 22Mab (ImClone Systems Incorporated of New York, New York, USA), the compounds ZD-1839 (AstraZeneca), BIBX-1382 (Boehringer Ingelheim), MDX-447 (Medarex Inc. of Annandale, New Jersey, USA), and OLX-103  
35 (Merck & Co. of Whitehouse Station, New Jersey, USA), VRCTC-310 (Ventech Research) and EGF fusion toxin (Seragen Inc. of Hopkinton, Massachusetts). These and other EGFR-inhibiting agents

can be used in the present invention.

VEGF inhibitors, for example SU-5416 and SU-6668 (Sugen Inc. of South San Francisco, California, USA), can also be combined with  
5 the compound of the present invention. VEGF inhibitors are described in, for example in WO 99/24440 (published May 20, 1999), PCT International Application PCT/IB99/00797 (filed May 3, 1999), in WO 95/21613 (published August 17, 1995), WO 99/61422 (published December 2, 1999), United States Patent 5,834,504  
10 (issued November 10, 1998), WO 98/50356 (published November 12, 1998), United States Patent 5,883,113 (issued March 16, 1999), United States Patent 5,886,020 (issued March 23, 1999), United States Patent 5,792,783 (issued August 11, 1998), WO 99/10349 (published March 4, 1999), WO 97/32856 (published September 12,  
15 1997), WO 97/22596 (published June 26, 1997), WO 98/54093 (published December 3, 1998), WO 98/02438 (published January 22, 1998), WO 99/16755 (published April 8, 1999), and WO 98/02437 (published January 22, 1998), all of which are incorporated herein in their entireties by reference. Other examples of some  
20 specific VEGF inhibitors useful in the present invention are IM862 (Cytran Inc. of Kirkland, Washington, USA); anti-VEGF monoclonal antibody of Genentech, Inc. of South San Francisco, California; and angiozyme, a synthetic ribozyme from Ribozyme (Boulder, Colorado) and Chiron (Emeryville, California). These  
25 and other VEGF inhibitors can be used in the present invention as described herein.

ErbB2 receptor inhibitors, such as GW-282974 (Glaxo Wellcome plc), and the monoclonal antibodies AR-209 (Aronex  
30 Pharmaceuticals Inc. of The Woodlands, Texas, USA) and 2B-1 (Chiron), can furthermore be combined with the compound of the invention, for example those indicated in WO 98/02434 (published January 22, 1998), WO 99/35146 (published July 15, 1999), WO 99/35132 (published July 15, 1999), WO 98/02437 (published  
35 January 22, 1998), WO 97/13760 (published April 17, 1997), WO

95/19970 (published July 27, 1995), United States Patent 5,587,458 (issued December 24, 1996), and United States Patent 5,877,305 (issued March 2, 1999), which are all hereby incorporated herein in their entireties by reference. ErbB2  
5 receptor inhibitors useful in the present invention are also described in United States Provisional Application No. 60/117,341, filed January 27, 1999, and in United States Provisional Application No. 60/117,346, filed January 27, 1999, both of which are incorporated in their entireties herein by  
10 reference. The erbB2 receptor inhibitor compounds and substance described in the aforementioned PCT applications, U.S. patents, and U.S. provisional applications, as well as other compounds and substances that inhibit the erbB2 receptor, can be used with the compound of the present invention in accordance with the  
15 present invention.

The compound of the invention can also be used with other agents useful in treating abnormal cell growth or cancer, including, but not limited to, agents capable of enhancing antitumor immune  
20 responses, such as CTLA4 (cytotoxic lymphocyte antigen 4) antibodies, and other agents capable of blocking CTLA4; and anti-proliferative agents such as farnesyl protein transferase inhibitors. Specific CTLA4 antibodies that can be used in the present invention include those described in United States  
25 Provisional Application 60/113,647 (filed December 23, 1998) which is incorporated by reference in its entirety, however other CTLA4 antibodies can be used in the present invention.

Such conjoint treatment may be achieved by way of the  
30 simultaneous, sequential or separate dosing of the individual components of the treatment.

It is expected that the compound of the invention can render abnormal cells more sensitive to treatment with radiation for  
35 purposes of killing and/or inhibiting the growth of such cells.



Accordingly, this invention further relates to a method for sensitizing abnormal cells in a mammal to treatment with radiation which comprises administering to the mammal an amount of the compound of the invention, pharmaceutically acceptable  
5 salt or solvate thereof, or prodrug thereof, which amount is effective in sensitizing abnormal cells to treatment with radiation. The amount of the compound, salt, solvate, or prodrug in this method can be determined according to the means for ascertaining effective amounts of the compound of the  
10 invention described herein.

The subject invention also includes isotopically-labelled compounds, which compounds are identical to the above recited compound of the invention, but for the fact that one or more  
15 atoms thereof are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into the compound of the invention include isotopes of hydrogen, carbon, nitrogen and oxygen, such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{13}\text{C}$ ,  
20  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{18}\text{O}$  and  $^{17}\text{O}$ , respectively. Compounds of the present invention, and pharmaceutically acceptable salts of said compounds which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present  
25 invention, for example those into which radioactive isotopes such as  $^3\text{H}$  and  $^{14}\text{C}$  are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e.,  $^3\text{H}$ , and carbon-14, i.e.,  $^{14}\text{C}$ , isotopes are particularly preferred for their ease of preparation and detectability. Further,  
30 substitution with heavier isotopes such as deuterium, i.e.,  $^2\text{H}$ , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of this  
35 invention can generally be prepared by carrying out the

procedures disclosed in the Methods and/or the examples below, and substituting a readily available isotopically labelled reagent for a non-isotopically labelled reagent, using methods well known in the art. Accordingly, reference to the compound 5 of the invention for use in the therapeutic methods and pharmaceutical compositions described herein also encompasses isotopically-labelled forms of the compound.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-  
10 amine hydrochloride, preferably the stable polymorph B form, is typically used at doses of 1-7000 mg/day, preferably 5-2500 mg/day, most preferably 5-200 mg/day, for any of the above treatments.

15 Patients that can be treated with the compound of the invention, alone or in combination, include, for example, patients that have been diagnosed as having psoriasis, BPH, lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head and neck, cutaneous or intraocular melanoma, ovarian cancer, rectal  
20 cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, gynecologic tumors (e.g., uterine sarcomas, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina or carcinoma of the vulva), Hodgkin's disease, cancer of the esophagus,  
25 cancer of the small intestine, cancer of the endocrine system (e.g., cancer of the thyroid, parathyroid or adrenal glands), sarcomas of soft tissues, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemia, solid tumors of childhood, lymphocytic lymphomas, cancer of the bladder,  
30 cancer of the kidney or ureter (e.g., renal cell carcinoma, carcinoma of the renal pelvis), or neoplasms of the central nervous system (e.g., primary CNS lymphoma, spinal axis tumors, brain stem gliomas or pituitary adenomas).

### Activity

The *in vitro* activity of the compounds of the present invention in inhibiting the receptor tyrosine kinase (and thus subsequent proliferative response, e.g., cancer) may be determined by the following procedure.

The activity of the compounds of the present invention, *in vitro*, can be determined by the amount of inhibition of the phosphorylation of an exogenous substrate (e.g., Lys<sub>3</sub> - Gastrin or polyGluTyr (4:1) random copolymer (I. Posner et al., J. Biol. Chem. 267 (29), 20638-47 (1992)) on tyrosine by epidermal growth factor receptor kinase by a test compound relative to a control. Affinity purified, soluble human EGF receptor (96 ng) is obtained according to the procedure in G. N. Gill, W. Weber, Methods in Enzymology 146, 82-88 (1987) from A431 cells (American Type Culture Collection, Rockville, MD) and preincubated in a microfuge tube with EGF (2µg/ml) in phosphorylation buffer + vanadate (PBV: 50 mM HEPES, pH 7.4; 125 mM NaCl; 24 mM MgCl<sub>2</sub>; 100 µM sodium orthovanadate), in a total volume of 10 µl, for 20-30 minutes at room temperature. The test compound, dissolved in dimethylsulfoxide (DMSO), is diluted in PBV, and 10 µl is mixed with the EGF receptor /EGF mix, and incubated for 10-30 minutes at 30°C. The phosphorylation reaction is initiated by addition of 20 µl <sup>33</sup>P-ATP/ substrate mix (120 µM Lys<sub>3</sub>-Gastrin (sequence in single letter code for amino acids, KKKGPWLEEEEEAYGWLDF), 50 mM Hepes pH 7.4, 40 µM ATP, 2 µCi γ-[<sup>33</sup>P]-ATP) to the EGFr/EGF mix and incubated for 20 minutes at room temperature. The reaction is stopped by addition of 10 µl stop solution (0.5 M EDTA, pH 8; 2mM ATP) and 6 µl 2N HCl. The tubes are centrifuged at 14,000 RPM, 4°C, for 10 minutes. 35 µl of supernatant from each tube is pipetted onto a 2.5 cm circle of Whatman P81 paper, bulk washed four times in 5% acetic acid, 1 liter per wash, and then air dried. This results in the binding of substrate to the paper with loss of free ATP on washing. The [<sup>33</sup>P] incorporated is measured by liquid

scintillation counting. Incorporation in the absence of substrate (e.g., lys<sub>3</sub>-gastrin) is subtracted from all values as a background and percent inhibition is calculated relative to controls without test compound present. Such assays, carried  
5 out with a range of doses of test compounds, allow the determination of an approximate IC<sub>50</sub> value for the *in vitro* inhibition of EGFR kinase activity.

Other methods for determining the activity of the compounds of  
10 the present invention are described in U.S. Patent No. 5,747,498, the disclosure of which is incorporated herein.

#### Pharmaceutical Compositions

The pharmaceutical composition may, for example and most  
15 preferably, be in a form suitable for oral administration as a tablet, capsule, pill, powder, sustained release formulations, solution, and suspension. Less preferred (with the mesylate form being the preferred form) are compositions for parenteral injection as a sterile solution, suspension or emulsion, for  
20 topical administration as an ointment or cream or for rectal administration as a suppository. The pharmaceutical composition may be in unit dosage forms suitable for single administration of precise dosages. The pharmaceutical composition will include a conventional pharmaceutical carrier or excipient and a  
25 compound according to the invention as an active ingredient. In addition, it may include other medicinal or pharmaceutical agents, carriers, adjuvants, etc.

Exemplary parenteral administration forms include solutions or  
30 suspensions of active compounds in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired.

Suitable pharmaceutical carriers include inert diluents or  
35 fillers, water and various organic solvents. The pharmaceutical

compositions may, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus for oral administration, tablets containing various excipients, such as citric acid may be employed together with various  
5 disintegrants such as starch, alginic acid and certain complex silicates and with binding agents such as sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may  
10 also be employed in soft and hard filled gelatin capsules. Preferred materials, therefor, include lactose or milk sugar and high molecular weight polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration the active compound therein may be combined with various sweetening  
15 or flavoring agents, coloring matters or dyes and, if desired, emulsifying agents or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin, or combinations thereof. Additionally, it is also possible to administer the compound of the invention topically and this may  
20 be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

The compound of the invention may also be administered to a mammal other than a human. The dosage to be administered to a  
25 mammal will depend on the animal species and the disease or disorder being treated. The compound may be administered to animals in the form of a capsule, bolus, tablet or liquid drench. The compound may also be administered to animals by injection or as an implant. Such formulations are prepared in  
30 a conventional manner in accordance with standard veterinary practice. As an alternative, the compound may be administered with the animal feedstuff, and for this purpose a concentrated feed additive or premix may be prepared for mixing with the normal animal feed.

Methods of preparing various pharmaceutical compositions with a specific amount of active compound are known, or will be apparent, to those skilled in this art. For examples, see Remington's Pharmaceutical Sciences, Mack Publishing Company, 5 Easter, Pa., 15th Edition (1975).

#### Administration and Dosage

Administration of the compounds of the present invention (hereinafter the "active compound(s)") can be effected by any  
10 method that enables delivery of the compounds to the site of action. These methods preferably include oral routes such as in the form of tablets, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion), topical, and rectal administration.  
15 While parenteral administration is usually preferred, oral administration is preferred for the hydrochloride B polymorph.

The amount of the active compound administered will be dependent on the subject being treated, the severity of the disorder or  
20 condition, the rate of administration and the judgement of the prescribing physician. However, an effective dosage is in the range of about 0.001 to about 100 mg per kg body weight per day, preferably about 1 to about 35 mg/kg/day, in single or divided doses. For a 70 kg human, this would amount to about 0.05 to  
25 about 7 g/day, preferably about 0.2 to about 2.5 g/day. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into  
30 several small doses for administration throughout the day.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride, preferably the stable polymorph B form, at doses of 1-7000 mg/day, preferably 5-2500 mg/day, most  
35 preferably 5-200 mg/day, is also useful for the treatment of patients (as measured, for example, by increased survival times)

by using combination therapies, for example in NSCLC (IIIb/V), as a 1<sup>st</sup> line therapy with carboplatin/paclitaxel or gemcitabine/cisplatin, in NSCLC (IIIb/V), as a 2<sup>nd</sup> line therapy with taxotere, and in head and neck cancers, as a 2nd line  
5 therapy with methotrexate for patients refractory to 5FU/cisplatin.

[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride, preferably the stable polymorph B form, at  
10 doses of 1-7000 mg/day, preferably 5-2500 mg/day, most preferably 5-200 mg/day, is also useful for the treatment of patients with additional conditions, including pancreatic cancer, with or without gemcitabine co-treatment, as first line therapy, for renal cancer, gastric cancer, prostate cancer,  
15 colorectal cancer (e.g. as a 2nd line therapy for patients who have failed 5FU/LCV/Irinotecan therapy), and also for hepatocellular, bladder, brain, ovarian, breast, and cervical cancers. For such treatments, in advanced disease patients with refractory disease, treatment effectiveness is readily monitored  
20 by an increased response rate, an increased time to progression or an increase in survival time.


[6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride, preferably the stable polymorph B form, is  
25 typically used at doses of 1-7000 mg/day, preferably 5-2500 mg/day, most preferably 5-200 mg/day, for any of the above treatments.

The examples and preparations provided below further illustrate  
30 and exemplify the compounds of the present invention and methods of preparing such compounds. This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the  
35 invention as described more fully in the claims which follow thereafter.

# EXPERIMENTAL DETAILS

## Example 1 - Preparation of compound of formula 4

### Reaction:



5 The following materials were used in the synthesis of the compound of formula 4:

Materials	Quantity	Units	Equivalents/Volumes
Compound of formula 3	88.0	kg	1 equivalent
10 Thionyl chloride	89.0	kg	2.5 equivalents
Dimethylformamide	11	kg	0.5 equivalent
methylene chloride	880.0	L	10 L/kg
50% sodium hydroxide solution	as required	L	1 equivalent
15 Heptane	880.0	L	10 L/kg

The following procedure is exemplary of the procedure to follow in the synthesis of the formula 4 compound:

88.0 kg of the compound of formula 3, 880.0 L methylene chloride, and 11.0 kg of dimethylformamide were charged to a clean, dry, glass-lined vessel under nitrogen atmosphere. 89 Kg of thionyl chloride were added to the mix while it is maintained at a temperature of a less than 30 °C during the charge. The contents of the reaction vessel were then heated for a minimum of five hours at reflux temperature before sampling for reaction completion and the pH is adjusted to be maintained between 7.0 to 8.0, by using 50 % NaOH, as required and the temperature of the reaction mixture is maintained at less than 25 °C. The biphasic mixture is stirred for fifteen to twenty minutes and allowed to settle for a minimum of thirty minutes. The layers were separated



and the organic layer was concentrated to 1/3 of its volume by removing methylene chloride. 880 L heptane was added with continued distillation of the remaining methylene chloride until the distillate reaches a temperature between 65 and 68 °C. The mixture was then cooled to between 10 to 15 °C over 5 hours and granulated for a minimum of 1 hour with the solids being isolated by filtration and washed with 220 L heptane. The solids (formula 4 compound) were dried in a vacuum drier at 45 to 50 °C.

#### Example 2 - Alternative Preparation of Compound of formula 4

In the reaction shown in Example 1, sodium bicarbonate may successfully be used instead of sodium hydroxide as shown in this Example.

Materials	Quantity	Units	Equivalents/Volumes
Compound of formula 3	30.0	kg	1 equivalent
Thionyl chloride	36.4	kg	3 equivalents
Dimethylformamide	3.75	kg	0.5 equivalent
methylene chloride	300	L	10 L/kg
50% sodium hydroxide solution	as required	L	
Heptane	375	L	12.5 L/kg
Heptane (wash)	90	L	3 L/kg
Sodium Bicarbonate	64.2	Kg	7.5 equivalents


30.0 kg of the compound of formula 3, 300.0L methylene chloride, and 3.75 kg of dimethylformamide were charged to a clean, dry, glass-lined vessel under a nitrogen atmosphere. 36.4 kg of thionyl chloride was added to the mix while it was maintained at a temperature of less than 30°C during the charge. The contents of the reaction vessel were then heated at reflux temperature for 13h before sampling for reaction completion. The reaction mixture was cooled to 20-25°C and added slowly to a stirred solution of sodium bicarbonate 64.2 kg and water 274L cooled to 4°C so that the temperature was maintained at less than 10°C. The final pH of the mixture was adjusted to within the range 7.0 to 8.0 by using 50% sodium hydroxide solution as required. The biphasic mixture

was stirred for fifteen to twenty minutes and allowed to settle for a minimum of thirty minutes at 10-20°C. The layers were separated and the organic layer was concentrated to 1/3 of its volume by removing methylene chloride. 375L of heptane was added with continued distillation of the remaining methylene chloride until the distillate reached a temperature between 65 and 68°C. The mixture was then cooled to 0 to 5°C over 4 hour and granulated for a minimum of 1 hour with the solids being isolated by filtration and washed with 90L heptane.

The solids (formula 4 compound) were dried in a vacuum drier at 45 to 50°C.

### 15 Example 3 -Preparation of compound of formulas 6 and 2 (Step 2):

#### Reaction:



The following materials were used in the synthesis of the compound of formula 6, as intermediate, and the compound of formula 2:

	Materials	Quantity	Units	Equivalents/Volumes
	Compound of formula 5	61.1	kg	1.2 equivalents
	Toluene	489	L	8 L/kg (WRT to formula 5 c'mpd)
25	Sodium hydroxide pellets	4.5	kg	0.16 equivalents
	Filteraid	0.5	kg	0.017 kg/kg (WRT to c'mpd 5)
	Compound of formula 4	90.8	kg	1.0 equivalent
	Acetonitrile	732	L	12 L/kg (WRT to c'mpd 5)

#### Example 4 - Preparation of compound of formula 2

The following procedure is exemplary of the procedure to follow in the synthesis of the formula 2 compound and intermediate compound of formula 6:

5

10

15

20

61.1 kg of formula 5 compound, 4.5 kg sodium hydroxide pellets and 489 L toluene were charged to a clean, dry, reaction vessel under nitrogen atmosphere and the reaction temperature is adjusted to between 105 to 108 °C. Acetone was removed over four hours by atmospheric distillation while toluene is added to maintain a minimum volume of 6 L of solvent per kg of formula 5 compound. The reaction mixture was then heated at reflux temperature, returning distillates to pot, until the reaction was complete. The mixture was then cooled to between 20 to 25 °C, at which time a slurry of 40.0 L toluene and 0.5 kg filteraid was charged to the reaction mixture and the mixture was agitated for ten to fifteen minutes. The resultant material was filtered to remove filteraid, and the cake is washed with 30 L toluene (compound of formula 6).

25

30

35


The filtrate (compound of formula 6) was placed in a clean, dry reaction vessel under nitrogen atmosphere, and 90.8 kg of the compound of formula 4 was charged into the reaction vessel together with 732 L acetonitrile. The reaction vessel was heated to reflux temperature and well agitated. Agitator speed was lowered when heavy solids appear. When the reaction was complete, the contents of reaction vessel were cooled to between 19 to 25 °C over three to four hours and the contents were agitated for at least one hour at a temperature between 20 and 25 °C. The solids (compound of formula 2, polymorph A form, or mixture of polymorph A and B) were then isolated by filtration and the filter cake was washed with two portions of 50 L acetonitrile and dried under vacuum at a temperature between 40 and 45 °C.

It has been discovered that the production of the A polymorph is favored by the reduction of the amount of acetonitrile relative to toluene, and particularly favored if isopropanol is used in place of acetonitrile. However, the use of isopropanol or other alcohols as cosolvents is disfavored because of the propensity to form an ether linkage between the alcoholic oxygen and the 4-carbon of the quinazoline, instead of the desired ethynyl phenyl amino moiety.

10 It has been further discovered that adjusting the pH of the reaction to between pH 1 and pH 7, preferably between pH 2 and pH 5, more preferably between pH 2.5 and pH 4, most preferably pH 3, will improve the rate of the reaction.

15 **Example 5 - Recrystallization of compound of formula 2 (which may be in polymorph A form or a mixture of polymorphs A and B) to Polymorph B (Step 3)**

**Reaction:**



The following materials were used in the conversion of polymorph A (or mixtures of polymorphs A and B) to polymorph B of the compound of formula 2:

	Materials	Quantity	Units	Equivalents/Volumes
30	Polymorph A (formula 2)	117.6	kg	1 equivalent
	2B-ethanol	1881.6	L	16 L/kg
	Water	470.4	L	4 L/kg

The following procedure is exemplary of procedures used to convert polymorph A (or mixtures of polymorphs A and B) into the more thermodynamically stable polymorph B of the compound of formula 2:

117.6 kg of the polymorph A (or mixtures of polymorphs A and B) were charged to a clean, dry, reaction vessel together 1881.6 L 2B-ethanol and 470.4 L water under a nitrogen atmosphere. The temperature was adjusted to reflux (~80 °C) and the mixture was agitated until the solids dissolve. The solution was cooled to between 65 and 70 °C and clarified by filtration. With low speed agitation, the solution was further cooled to between 50 and 60 °C over a minimum time of 2 hours and the precipitate was granulated for 2 hours at this temperature. The mixture was further cooled to between 0 and 5 °C over a minimum time of 4 hours and granulated for a minimum of 2 hours at this temperature. The solids (polymorph B) were isolated by filtration and washed with at least 100 L 2B-ethanol. The solids were determined to be crystalline polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride substantially free of the polymorph A from. The solids obtained by this method are substantially homogeneous polymorph B form crystals relative to the polymorph A form. The method allows for production of polymorph B in an amount at least 70% by weight, at least 80% by weight, at least 90% by weight, at least 95% by weight, and at least 98% by weight relative to the weight of the polymorph A. It is to be understood that the methods described herein are only exemplary and are not intended to exclude variations in the above parameters which allow the production of polymorph B in varying granulations and yields, according to the desired storage, handling and manufacturing applications of the compound. The solids were vacuum dried at a temperature below 50°C and the resultant product was milled to provide the polymorph B in usable form.

**Example 6 - Clinical studies utilizing treatment with the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride.**

5 The stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride is a potent, selective and orally active inhibitor of the epidermal growth factor receptor (EGFR) protein-tyrosine kinase, an oncogene that has been associated with the aberrant  
10 growth that is characteristic of cancer cells. This compound is being evaluated in clinical trials in normal healthy volunteers and in cancer patients in order to assess its safety profile and effectiveness.

15 Phase I Clinical Studies

Phase I clinical studies of the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride have been effectively completed in volunteers, initially, and subsequently in cancer patients, at single doses  
20 ranging from 25-200 mg/day or 100-1600 mg/week. Data from these studies revealed no adverse events that were greater than moderate in severity for a dose of 150 mg/day. In a daily dosing regimen study the dose limiting toxicity at 200 mg/day was diarrhea. This observed side effect was effectively controlled  
25 at the 150mg daily dose level using Loperamide (Imodium®). The second adverse event observed in these studies, and most significant toxicity at 150 mg daily, was a monomorphic acneiform rash analogous to that reported for other EGFR inhibitor agents in clinical trials. This rash had an "above-  
30 waist" distribution including face, scalp, neck, arms, chest and back. The rash has a unique histopathology of PMN infiltration with mild epidermal hyperproliferation. It is not consistent with drug hypersensitivity nor does it appear to be a "named" dermatological condition. This rash has not been a significant  
35 impediment to patients staying on the Phase II trials. The stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-

4-yl]-(3-ethynylphenyl)-amine hydrochloride has been tested in a total of 290 patients in Phase I and ongoing Phase II studies and demonstrates a well tolerated safety profile. Furthermore, preliminary evidence of effectiveness was observed in Phase I studies. For example, in one Phase I study of 28 patients, 8 patients remain alive over a year after inception of treatment and 12 patients remained alive from 9-22 months.

In order to establish a suitable safety profile, the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride is also used at doses of 1-7000 mg/day, preferably 5-2500 mg/day, most preferably 5-200 mg/day, in Phase I clinical combination studies with one or more additional drugs or treatments, preferably selected from one of the following group - Taxol, Gemcitabine, Taxotere, Capcitabine, 5FU, Cisplatin, Temozolomide, radiation treatment, and chemoradiation treatment.

#### Phase II and Phase III Clinical Studies

Three Phase II single agent studies of the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride in refractory non-small cell lung cancer, advanced head and neck cancer and refractory ovarian cancer, at a 150 mg daily dose were initiated.

Indications of single agent anti-tumor activity for the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride was seen in patients with advanced cancers in several different tumor types. For example, initial findings indicate that the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride is a well-tolerated oral medication that is active as a monotherapy when administered to patients with advanced head and neck cancer. In preliminary results 3 patients had objective partial responses, while another 9 patients showed evidence of a stabilization of their disease status. The

acneiform rash, which is apparently characteristic of all the anti-EGFR inhibitors undergoing clinical testing, was reported in approximately 70% of the first group of patients in this study.

5

The early data emerging from the 48 patient Phase II study in refractory non-small cell lung cancer (NSCLC) patients also indicates the effectiveness of treatment with the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-  
10 (3-ethynylphenyl)-amine hydrochloride as a single agent anti-tumor drug for NSCLC. Of the first 19 evaluable patients in the study, 5 had objective partial responses, while another 4 patients showed evidence of a stabilization of their disease status. Partial responses were observed in two patients who had  
15 been treated previously with two and three different chemotherapy regimens. Thus it appears that the stable polymorph B form of [6,7-bis(2-methoxyethoxy)quinazolin-4-yl]-(3-ethynylphenyl)-amine hydrochloride is a well tolerated, oral medication which is active in non-small cell lung cancer.

20

Qualification criteria for the open label, single agent study required the patients to have failed platinum-based chemotherapy and to have tumors that are histopathologically confirmed to be EGFR positive. The primary endpoint in the study is response  
25 rate with stable disease and time-to-progression amongst the secondary end-points.

Evidence of anti-tumor activity can also be seen in the patients with ovarian cancer in the on-going Phase II study. In  
30 preliminary results 2 patients had objective partial responses, while another 4 patients showed evidence of a stabilization of their disease status. Documented evidence of anti-tumor activity was also seen in other EGFR positive tumor types, including colorectal and renal cell carcinoma, from Phase I studies in  
35 cancer patients with multiple tumor types.



What is claimed is:

1. A substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91.
2. The polymorph of claim 1, characterized by the X-ray powder diffraction pattern shown in Figure 3.
3. A crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91, which is substantially free of the A polymorph.
4. The polymorph of claim 3, characterized by the X-ray powder diffraction pattern shown in Figure 3.
5. A composition comprising a substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91.

6. The composition of claim 5, wherein the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately:

2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)
6.255	100.0	17.668	2.5	22.982	4.8	27.534	0.9	32.652	1.7
7.860	3.2	18.193	0.7	23.589	2.3	28.148	1.5	33.245	1.7
9.553	3.9	18.749	1.5	23.906	3.0	28.617	4.3	34.719	1.5
11.414	1.5	19.379	1.0	24.459	6.8	29.000	1.4	35.737	0.8
12.483	6.4	20.196	14.4	25.138	10.0	29.797	2.1	36.288	1.0
13.385	9.6	20.734	4.2	25.617	3.7	30.267	0.9	36.809	0.6
14.781	2.1	21.103	14.4	25.908	3.9	30.900	1.6	37.269	1.1
15.720	2.9	21.873	4.7	26.527	2.8	31.475	2.2	37.643	1.4
16.959	5.5	22.452	4.5	26.911	5.6	31.815	2.4	38.114	1.7

7. The composition of claim 5, wherein the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the polymorph B form is characterized by the X-ray powder diffraction pattern shown in Figure 3.
8. A composition comprising a crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91 in a weight % of the B polymorph relative to the A polymorph which is at least 70%.
9. The composition of claim 8, wherein the B polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine exhibits an X-ray powder diffraction pattern having characteristic peaks expressed


in degrees 2-theta at approximately:

2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)
6.255	100.0	17.668	2.5	22.982	4.8	27.534	0.9	32.652	1.7
7.860	3.2	18.193	0.7	23.589	2.3	28.148	1.5	33.245	1.7
9.553	3.9	18.749	1.5	23.906	3.0	28.617	4.3	34.719	1.5
11.414	1.5	19.379	1.0	24.459	6.8	29.000	1.4	35.737	0.8
12.483	6.4	20.196	14.4	25.138	10.0	29.797	2.1	36.288	1.0
13.385	9.6	20.734	4.2	25.617	3.7	30.267	0.9	36.809	0.6
14.781	2.1	21.103	14.4	25.908	3.9	30.900	1.6	37.269	1.1
15.720	2.9	21.873	4.7	26.527	2.8	31.475	2.2	37.643	1.4
16.959	5.5	22.452	4.5	26.911	5.6	31.815	2.4	38.114	1.7


10. The composition of claim 8, wherein the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the polymorph B form is characterized by the X-ray powder diffraction pattern shown in Figure 3.
11. A pharmaceutical composition which comprises a therapeutically effective amount of the polymorph of claim 1 and a pharmaceutically acceptable carrier.
12. The pharmaceutical composition of claim 11, wherein said composition is adapted for oral administration.
13. The pharmaceutical composition of claim 12, wherein the pharmaceutical composition is in the form of a tablet.
14. A method of treating a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically effective amount of the polymorph of claim 1.
15. The method of claim 14, wherein the method is for the treatment of a cancer selected from brain, squamous cell, bladder, gastric, pancreatic, breast, head, neck, oesophageal, prostate, colorectal, lung, renal, kidney, ovarian, gynecological and thyroid cancer.

16. The method of claim 14, wherein the method of for the treatment of a cancer selected from non-small cell lung cancer (NSCLC), refractory ovarian cancer, head and neck cancer, colorectal cancer and renal cancer.
17. The method of claim 14, wherein the therapeutically effective amount is from about 0.001 to about 100 mg/kg/day.
18. The method of claim 14, wherein the therapeutically effective amount is from about 1 to about 35 mg/kg/day.
19. The method of claim 14, wherein the therapeutically effective amount is from about 1 to about 7000 mg/day.
20. The method of claim 19, wherein the therapeutically effective amount is from about 5 to about 2500 mg/day.
21. The method of claim 20, wherein the therapeutically effective amount is from about 5 to about 200 mg/day.
22. The method of claim 21, wherein the therapeutically effective amount is from about 25 to about 200 mg/day.
23. A method for the treatment of a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically effective amount of the polymorph of claim 1 in combination with an anti-tumor agent selected from the group consisting of a mitotic inhibitor, an alkylating agent, an anti-metabolite, an intercalating antibiotic, a growth factor inhibitor, a cell cycle inhibitor, an enzyme, a topoisomerase inhibitor, a biological response modifier, an anti-hormone, and an anti-androgen.


24. A method of preparing a crystalline polymorph of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride designated the B polymorph which comprises the step of recrystallizing N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in a solvent comprising alcohol.
25. The method of claim 24, wherein the solvent further comprises water.
26. The method of claim 24, wherein N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride is prepared by coupling a compound of formula 6



with a compound of formula 4




27. The method of claim 26, wherein said compound of formula 6 is prepared by reacting a compound of formula 5



in a suspension of metal alkali and solvent and with heating.


28. The method of claim 26, wherein said compound of formula 4 is prepared by chlorinating a compound of formula 3



3.


29. A method for the production of the polymorph B of claim 1 comprising the steps of:

a) substitution chlorination of starting quinazolinamine compound of formula 3



3


having an hydroxyl group, to provide a compound of formula 4



4


by reaction thereof in a solvent mixture of thionyl chloride, methylene chloride and dimethylformamide,

b) preparation of a compound of formula 6



6

in situ from starting material of compound of formula 5



5

by reaction of the latter in a suspension of metal alkali and solvent and with heating;

c) reaction of the compound of formula 6 in situ with the compound of formula 4 wherein the compound of formula 6 replaces the chlorine in the compound of formula 4 to give the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride;

d) recrystallizing the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride, in alcohol, into the polymorph B form.

30. The method of claim 29, wherein the substitution chlorination is quenched in the presence of aqueous sodium hydroxide.

31. The method of claim 29, wherein the substitution chlorination is quenched in the presence of aqueous sodium bicarbonate.
32. The method of claim 29, wherein the substitution chlorination is quenched in the presence of aqueous potassium hydroxide, aqueous potassium bicarbonate, aqueous potassium carbonate, aqueous sodium carbonate, or a mixture thereof.
33. A method for the production of a crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph by recrystallization comprising the steps of:
  - e) heating to reflux alcohol, water and the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine so as to form a solution;
  - f) cooling the solution to between about 65 and 70 °C;
  - g) clarifying the solution; and
  - h) precipitating polymorph B by further cooling the clarified solution.
34. A substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the A polymorph characterized by the X-ray powder diffraction pattern shown in Figure 1.
35. A crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the A polymorph characterized by the X-ray powder diffraction pattern shown in Figure 1, which is substantially free of the B polymorph.



36. A composition comprising a substantially homogeneous crystalline polymorph of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride designated the A polymorph, which is characterized by the following peaks in its X-ray powder diffraction pattern shown in Figure 1.

37. A composition comprising a substantially homogeneous crystalline polymorph of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the form of polymorph A, which is characterized by the following peaks:

Polymorph A

Anode: Cu- Wavelength 1: 1.54056 Wavelength 2: 1.54439 (Rel Intensity: 0.500)

Range# 1 - Coupled: 3.000 to 40.000 StepSize: 0.040 StepTime: 1.00

Smoothing Width: 0.300 Threshold: 1.0

d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)
15.82794	100.0	6.63179	1.7	4.54453	4.8	3.61674	8.2	2.91238	3.5
14.32371	3.9	5.84901	2.1	4.19685	4.7	3.50393	9.3	2.73148	3.7
11.74376	1.5	5.69971	2.3	4.16411	4.4	3.40200	6.0	2.60193	1.8
11.03408	1.2	5.46922	2.4	3.97273	4.7	3.35174	5.3	2.48243	1.3
10.16026	1.4	5.21396	3.6	3.91344	12.4	3.29005	4.2	2.40227	2.2
8.98039	13.1	4.80569	3.5	3.78223	24.2	3.05178	7.1	2.31297	1.7
7.85825	7.8	4.70077	12.2	3.67845	8.8	2.97750	3.0		

or,

Polymorph A

Anode: Cu - Wavelength 1: 1.54056 Wavelength 2: 1.54439 (Rel Intensity: 0.500)

Range#1 - Coupled: 3.000 to 40.000 StepSize: 0.040 StepTime: 1.00

Smoothing Width: 0.300 Threshold: 1.0

2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)
5.579	100.0	13.340	1.7	19.517	4.8	24.594	8.2	30.673	3.5
6.165	3.9	15.135	2.1	21.152	4.7	25.398	9.3	32.759	3.7
7.522	1.5	15.534	2.3	21.320	4.4	26.173	6.0	34.440	1.8
8.006	1.2	16.193	2.4	22.360	4.7	26.572	5.3	36.154	1.3
8.696	1.4	16.991	3.6	22.703	12.4	27.080	4.2	37.404	2.2
9.841	13.1	18.447	3.5	23.502	24.2	29.240	7.1	38.905	1.7
11.251	7.8	18.862	12.2	24.175	8.8	30.007	3.0		

38. A composition comprising a substantially homogeneous crystalline polymorph of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the form of polymorph B, which is characterized by the following peaks:

Polymorph B

Anode: Cu - Wavelength 1 1.54056 Wavelength 2: 1.54439 (Rel Intensity:0.500)

Range # 1 - Coupled 3.000 to 40.040 StepSize: 0.040 StepTime 1.00

Smoothing Width: 0.300 Threshold: 1.0

d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)	d(A)	I(rel)
14.11826	100.0	5.01567	2.5	3.86656	4.8	3.23688	0.9	2.74020	1.7
11.23947	3.2	4.87215	0.7	3.76849	2.3	3.16755	1.5	2.69265	1.7
9.25019	3.9	4.72882	1.5	3.71927	3.0	3.11673	4.3	2.58169	1.5
7.74623	1.5	4.57666	1.0	3.63632	6.8	3.07644	1.4	2.51043	0.8
7.08519	6.4	4.39330	14.4	3.53967	10.0	2.99596	2.1	2.47356	1.0
6.60941	9.6	4.28038	4.2	3.47448	3.7	2.95049	0.9	2.43974	0.6
5.98828	2.1	4.20645	14.4	3.43610	3.9	2.89151	1.6	2.41068	1.1
5.63253	2.9	4.06007	4.7	3.35732	2.8	2.83992	2.2	2.38755	1.4
5.22369	5.5	3.95667	4.5	3.31029	5.6	2.81037	2.4	2.35914	1.7

or,

Polymorph B

Anode: Cu - Wavelength 1 1.54056 Wavelength 2: 1.54439 (Rel Intensity:0.500)

Range# 1 - Coupled: 3.000 to 40.040 StepSize 0.040 StepTime: 1.00

Smoothing Width:0.300 Threshold: 1.0

2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)	2-Theta	I(rel)
6.255	100.0	17.668	2.5	22.982	4.8	27.534	0.9	32.652	1.7
7.860	3.2	18.193	0.7	23.589	2.3	28.148	1.5	33.245	1.7
9.553	3.9	18.749	1.5	23.906	3.0	28.617	4.3	34.719	1.5
11.414	1.5	19.379	1.0	24.459	6.8	29.000	1.4	35.737	0.8
12.483	6.4	20.196	14.4	25.138	10.0	29.797	2.1	36.288	1.0
13.385	9.6	20.734	4.2	25.617	3.7	30.267	0.9	36.809	0.6
14.781	2.1	21.103	14.4	25.908	3.9	30.900	1.6	37.269	1.1
15.720	2.9	21.873	4.7	26.527	2.8	31.475	2.2	37.643	1.4
16.959	5.5	22.452	4.5	26.911	5.6	31.815	2.4	38.114	1.7

39. A prodrug of the compound of claim 1.
40. A method of inducing differentiation of tumor cells in a tumor comprising contacting the cells with an effective amount of the compound of claim 1, or a composition of claims 3 or 6 so as to thereby differentiate the tumor cells.

41. A method for the treatment of NSCLC (non small cell lung cancer), pediatric malignancies, cervical and other tumors caused or promoted by human papilloma virus (HPV), melanoma, Barrett's esophagus (pre-malignant syndrome), adrenal and skin cancers and auto immune, neoplastic cutaneous diseases and atherosclerosis in a mammal comprising administering to said mammal a therapeutically effective amount of a pharmaceutical composition comprised of at least one of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and pharmaceutically acceptable salts thereof in anhydrous and hydrate forms.
42. The method of claim 41, wherein the treatment further comprises a palliative or neo-adjuvant/adjuvant monotherapy.
43. The method of claim 41, wherein the treatment further comprises blocking epidermal growth factor receptors (EGFR).
44. The method of claim 41, for use in treatment of tumors that express EGFRvIII.
45. The method of claim 41, wherein the treatment further comprises a combination with any of chemotherapy and immunotherapy
46. The method of claim 41, wherein the treatment further comprises, treatment with either or both anti-EGFR and anti-EGF antibodies.
47. The method of claim 41, wherein the treatment further comprises a further administration to said mammal of a member of the group consisting of inhibitors of MMP (matrix-metallo-proteinase), VEGFR (vascular endothelial growth factor receptor), farnesyl transferase, CTLA<sub>4</sub> (cytotoxic T-lymphocyte antigen 4) and erbB2, MAb to VEGFr, rhuMAb-VEGF, erbB2 MAb and avb3 Mab.

48. The method of claim 41, wherein the pharmaceutical compounds are used as radiation sensitizers for cancer treatment or in combination with anti-hormonal therapies.
49. The method of claim 41, wherein the pharmaceutical compounds are used for the inhibition of tumor growth in humans in a regimen with radiation treatment.
50. A method for the chemoprevention of basal or squamous cell carcinoma of the skin in areas exposed to the sun or in persons of high risk to said carcinoma, said method comprising administering to said persons a therapeutically effective amount of a pharmaceutical composition comprised of at least one of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and pharmaceutically acceptable salts thereof in anhydrous and hydrate forms.
51. A method of inducing differentiation of tumor cells in a tumor comprising contacting the cells with an effective amount of the compound of at least one of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, and pharmaceutically acceptable salts thereof in anhydrous and hydrate forms.
52. A method of making a composition which composition comprises substantially homogeneous crystalline polymorph of the hydrochloride salt of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine designated the B polymorph that exhibits an X-ray powder diffraction pattern having characteristic peaks expressed in degrees 2-theta at approximately 6.26, 12.48, 13.39, 16.96, 20.20, 21.10, 22.98, 24.46, 25.14 and, 26.91, comprising admixing the crystalline polymorph of claim 1 with a carrier.

53. The method of claim 52, wherein the N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride in the polymorph B form is characterized by the X-ray powder diffraction pattern shown in Figure 3.
54. The method of claim 52, wherein the carrier is a pharmaceutically acceptable carrier.

**STABLE POLYMORPH OF  
N-(3-ETHYNYLPHENYLAMINO)-6,7-BIS(2-  
METHOXYETHOXY)-4-QUINAZOLINAMINE HYDROCHLORIDE,  
METHODS OF PRODUCTION, AND PHARMACEUTICAL USES THEREOF**

Abstract of the Invention

The present invention relates to a stable crystalline form of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine hydrochloride designated the B polymorph, its production in essentially pure form, and its use. The invention also relates to the pharmaceutical compositions containing the stable polymorph B form of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine as hydrochloride, as well other forms of the compound, and to methods of treating hyperproliferative disorders, such as cancer, by administering the compound.

CP-358774-1 35076-183-1 - NORRIS

Figure 1

CP-358774-1 35076-183-1 - NORRIS





Figure 2

CP-358774-1 35076-183-1 - NORRIS






CP-358774-1 35076-076-1 - NORRIS

Figure 3


CP-358774-1 35076-076-1 - NORRIS



CP-358774-1 35076-076-1  
2013-09-10 10:00:00  
CP-358774-1 35076-076-1

Figure 4

CP-358774-1 35076-076-1 - NORRIS



**DECLARATION AND POWER OF ATTORNEY**

***As a below-named inventor, I hereby declare that:***

*My residence, post office address, and citizenship are as stated below next to my name.*

*I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:*

STABLE POLYMORPH OF N-(3-ETHYNYLPHENYLAMINO)-6, 7-BIS(2-METHOXYETHOXY)-4-QUINAZOLINAMINE HYDROCHLORIDE, METHODS OF PRODUCTION, AND PHARMACEUTICAL USES THEREOF

*the specification of which:  
(check one)*

X is attached hereto

was filed on \_\_\_\_\_ as \_\_\_\_\_

Application Serial No. \_\_\_\_\_

and was amended \_\_\_\_\_ (if applicable)

***I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.***

**I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.**

*I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 (a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International Application which designated at least one country other than the United States, listed below. I have also identified below any foreign application for patent or inventor's certificate, or PCT International Application having a filing date before that of the earliest application from which priority is claimed:*

**Prior Foreign Application(s)**

**Priority Claimed**

[illegible]

**Declaration and Power of Attorney**

Page 2

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

<u>Provisional Application No.</u>	<u>Filing Date</u>	<u>Status</u>
<u>60/206,420</u>	<u>May 23, 2000</u>	<u>pending as of November 9, 2000</u>
<u>60/193,191</u>	<u>March 30, 2000</u>	<u>pending as of November 9, 2000</u>
<u>60/164,907</u>	<u>November 11, 1999</u>	<u>pending as of November 9, 2000</u>
<u>                    </u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States Application(s), or Section 365(c) of any PCT International Application(s) designating the United States listed below. Insofar as this application discloses and claims subject matter in addition to that disclosed in any such prior Application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56, which became available between the filing date(s) of such prior Application(s) and the national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
<u>N/A</u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>
<u>                    </u>	<u>                    </u>	<u>                    </u>

**And I hereby appoint**

John P. White (Reg. No. 28,678); Christopher C. Dunham (Reg. No. 22,031); Norman H. Zivin (Reg. No. 25,385); Jay H. Maioli (Reg. No. 27,213); William E. Pelton (Reg. No. 25,702); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Wendy E. Miller (Reg. No. 35,615); Richard S. Milner (Reg. No. 33,970); Robert T. Maldonado (Reg. 38,232); Paul Teng (40,837); Richard F. Jaworski (Reg. No. 33,515); Elizabeth M. Wieckowski (Reg. No. 42,226); Pedro C. Fernandez (Reg. No. 41,741); Gary J. Gershik (Reg. No. 39,992); Jane M. Love (Reg. No. 42,812); Spencer H. Schneider (Reg. No. 45,923) and Raymond A. Diperna (Reg. No. 44,063).

and each of them, all c/o Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, New York 10036, my attorneys, each with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent, to transact all business in the Patent and Trademark Office connected therewith and to file any International Applications which are based thereon under the provisions of the Patent Cooperation Treaty.

Please address all communications, and direct all telephone calls, regarding this application to:

John P. White \_\_\_\_\_ Reg. No. 28,678

Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, New York 10036  
Tel. (212) 278-0400

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or  
first joint inventor \_\_\_\_\_ Timothy Norris \_\_\_\_\_

Inventor's signature \_\_\_\_\_

Citizenship \_\_\_\_\_ Great Britian \_\_\_\_\_ Date of signature \_\_\_\_\_

Residence \_\_\_\_\_ 27 Friar Tuck Drive, Gales Ferry, CT 06335 \_\_\_\_\_

Post Office Address \_\_\_\_\_ same as residence address above \_\_\_\_\_

Full name of joint  
inventor (if any) \_\_\_\_\_ Jeffrey W. Raggon \_\_\_\_\_

Inventor's signature \_\_\_\_\_

Citizenship \_\_\_\_\_ United States of America \_\_\_\_\_ Date of signature \_\_\_\_\_

Residence \_\_\_\_\_ 754 Pendleton Road, North Stonington, CT 06359 \_\_\_\_\_

Post Office Address \_\_\_\_\_ same as residence address above \_\_\_\_\_

Full name of joint  
inventor (if any) \_\_\_\_\_ Richard D. Connell \_\_\_\_\_

Inventor's signature \_\_\_\_\_

Citizenship \_\_\_\_\_ United States of America \_\_\_\_\_ Date of signature \_\_\_\_\_

Residence \_\_\_\_\_ 23 Jeremy Drive, East Lyme, CT 06333 \_\_\_\_\_

Post Office Address \_\_\_\_\_ same as residence address above \_\_\_\_\_

Full name of joint inventor (if any) James D. Moyer

Inventor's signature \_\_\_\_\_

Citizenship United States of America Date of signature \_\_\_\_\_

Residence 5 Jefferson Drive, East Lyme, CT 06333

Post Office Address same as residence address above

\_\_\_\_\_

Full name of joint inventor (if any) Michael J. Morin

Inventor's signature \_\_\_\_\_

Citizenship United States of America Date of signature \_\_\_\_\_

Residence 8 Goundry Drive, Waterford, CT 06385

Post Office Address same as residence address above

\_\_\_\_\_

Full name of joint inventor (if any) Shama M. Kajiji

Inventor's signature \_\_\_\_\_

Citizenship United States of America Date of signature \_\_\_\_\_

Residence 190 Mistuxet Avenue, Mystic, CT 06355

Post Office Address same as residence address above

\_\_\_\_\_

Full name of joint inventor (if any) Barbara A. Foster

Inventor's signature \_\_\_\_\_

Citizenship United States of America Date of signature \_\_\_\_\_

Residence 31 Rossie Street, Mystic, CT 06355

Post Office Address same as residence address above

\_\_\_\_\_

Full name of joint inventor (if any) Karen J. Ferrante

Inventor's signature \_\_\_\_\_

Citizenship United States of America Date of signature \_\_\_\_\_

Residence 150 Adirondack Drive, East Greenwich, RI 02818

Post Office Address same as residence address above  
\_\_\_\_\_

Full name of joint inventor (if any) Sandra L. Silberman

Inventor's signature \_\_\_\_\_

Citizenship United States of America Date of signature \_\_\_\_\_

Residence 10 Heather Lane, Randolph, NJ 07869

Post Office Address same as residence address above  
\_\_\_\_\_

Full name of joint inventor (if any) \_\_\_\_\_

Inventor's signature \_\_\_\_\_

Citizenship \_\_\_\_\_ Date of signature \_\_\_\_\_

Residence \_\_\_\_\_

Post Office Address \_\_\_\_\_  
\_\_\_\_\_